


# Labimycin<sup>®</sup> LA

Oxytetracycline 300 mg/ml



## Long-acting and lasting



**Indicated for  
cattle & sheep.  
Now also  
for pigs!**



### Labimycin<sup>®</sup> LA

Longer action<sup>1</sup>  
for a time-dependent antibiotic<sup>2</sup>

**LABIANA**  
*always works*



## Broad-spectrum, long-acting category D antibiotic with a single application

*Histophilus somni* - *Mannheimia haemolytica* - *Pasteurella multocida* - *Mycoplasma bovis*  
*Arcanobacterium pyogenes* - *Dichelobacter nodosus* - *Fusobacterium necrophorum*  
*Prevotella melaninogenica* - *Chlamydophila* spp. - *Dermatophilus congolensis* - *Bordetella bronchiseptica*  
*Pasteurella* spp. - *Staphylococcus aureus* - *Streptococcus agalactiae* - *Escherichia coli* - *Chlamydia abortus*  
*Chlamydia psittaci* - *Mycoplasma* spp. - *Erysipelothrix rhusiopathiae*

### EACH ML CONTAINS

Oxytetracycline (as Oxytetracycline dihydrate) ..... 300 mg

### INDICATIONS

This veterinary medicinal product is indicated for the control and treatment of a wide range of common systemic, respiratory, urinary and local infections:

#### Cattle:

Treatment of respiratory infections caused by strains of *Histophilus somni*, *Mannheimia haemolytica*, *Pasteurella multocida* and *Mycoplasma bovis*.

Treatment of metritis caused by strains of *Arcanobacterium pyogenes*.

Treatment of navel/joint infections caused by strains of *Dichelobacter nodosus*, *Fusobacterium necrophorum* and *Prevotella melaninogenica*.

#### Sheep:

Treatment of respiratory infections caused by strains of *Mannheimia haemolytica* and *Pasteurella*.

Treatment of genital infections caused by strains of *Arcanobacterium pyogenes*, *Chlamydophila* spp. and *Dermatophilus congolensis*.

Treatment of navel/joint infections caused by strains of *Dichelobacter nodosus*, *Fusobacterium necrophorum* and *Prevotella melaninogenica*.

#### Pigs:

For the treatment of respiratory infections caused by strains of *Bordetella bronchiseptica* and *Pasteurella*.

For the treatment of Erysipelas caused by strains of *Erysipelothrix rhusiopathiae*.

For the treatment of Atrophic rhinitis caused by strains of *Bordetella bronchiseptica* and *Pasteurella multocida*.

#### Other infections:

Treatment of mastitis caused by strains of *Staphylococcus aureus*, *Streptococcus agalactiae* and *Escherichia coli*.

Treatment of enzootic abortion caused by strains of *Chlamydia abortus* and *Chlamydia psittaci*.

Treatment of genital infections and Poliartitis caused by strains of *Chlamydia abortus* and *Mycoplasma* spp.

### DOSAGE AND ADMINISTRATION ROUTE

Intramuscular use.

The general recommended dosage for a prolonged duration of activity of 5 to 6 days is a single deep intramuscular injection of 30 mg oxytetracycline/kg bodyweight (equivalent to 1 ml of the veterinary medicinal product/10 kg of bodyweight).

Maximum recommended dose at one site:

**Cattle:** ..... 15 ml

**Sheep:** ..... 5 ml

**Pigs:** ..... 10 ml

**Piglets 1 day:** ..... 0.2 ml

**Piglets 7 days:** ..... 0.3 ml

**Piglets 14 days:** ..... 0.4 ml

**Piglets 21 days:** ..... 0.5 ml

**Piglets over 21 days:** ..... 1 ml/10 kg

### WITHDRAWAL PERIODS

**Cattle: Meat and offal:** 35 days. **Milk:** 7 days.

**Sheep: Meat and offal:** 35 days. **Milk:** 9 days.

**Pigs: Meat and offal:** 28 days.

### CONTRAINDICATIONS

Do not use in cases of hypersensitivity to the active substance(s) or to any of the excipient(s).

### SPECIAL PRECAUTIONS FOR USE

Do not dilute this veterinary medicinal product.

If administered simultaneously with other treatments, use a separate injection site.

Before any infectious process, bacteriological confirmation of the diagnosis and a sensitivity test of the bacteria causing the process is recommended.

Use of the product should be based on identification and susceptibility testing of the target pathogen(s). If this is not possible, therapy should be based on epidemiological information and knowledge of susceptibility of the target pathogens at farm level, or at local/regional level.

Use of the product should be in accordance with official, national and regional antimicrobial policies.

### PRESENTATION

Amber glass vials.

100 ml and 250 ml vials.

Registry nº 4074 ESP.

Medication subject to veterinary prescription.

Administration under control or supervision of the veterinary.

### References:

<sup>1</sup> Labimycin® 300 mg/ml SPC.

<sup>2</sup> Vicente D, Pérez-Trallero E. Tetraciclinas, sulfamidas y metronidazol [Tetracyclines, sulfonamides, and metronidazole]. *Enferm Infecc Microbiol Clin*. 2010 Feb;28(2):122-30. Spanish. doi:10.1016/j.eimc.2009.10.002. PMID: 20097452.