



Mg x2



Double quantity
of magnesium



Calcio Inyectable Labiana

Injectable Solution



It combines twice the amount of magnesium and calcium with one injection

Clinical signs of hypocalcaemia

- ✓ Lack of appetite
- ✓ Dystocia
- ✓ Lack of balance
- ✓ Down cow
- ✓ Tympanism

Clinical signs of hypomagnesemia

- ✓ Irritability
- ✓ Head and ears upright
- ✓ Difficulty with movement
- ✓ Hyperventilation
- ✓ Aggressiveness

0 days withdrawal period in milk and meat

 **LABIANA**
siempre eficaz




Calcio Inyectable Labiana

Injectable solution



Treatment of physiological and pathological states that cause hypocalcaemia and hypomagnesemia



COMPOSITION PER ML

Calcium gluconate for injectable 216,18 mg (equivalent to 19,32 mg of calcium)
Magnesium chloride hexahydrate 60 mg (equivalent to 7,17 mg of magnesium)
Magnesium hypophosphate hexahydrate 51 mg (equivalent to 4,73 mg of magnesium)

INDICATIONS

Treatment of physiological and pathological states that cause hypocalcaemia accompanied by hypomagnesemia.

DOSAGE AND ROUTE OF ADMINISTRATION

Young bovine: 25-100 ml/animal.

Adult bovine: 250-310 ml/animal.

In bovine, for the animals that do not recover within 4-8 hours treatment should be reassessed and, if necessary, should be repeated.

Adult horse: 250-600 ml/animal.

Young horse: 25 - 100 ml/animal.

Sheep, goats and swine: 25-100 ml/animal.

Piglets: 3-5 ml/animal.

Bibliography:

- Hilton WM. Nutritional Requirements. In: The Merck Veterinary Manual. 11th ed. Kenilworth: Merck & Co; 2016.
- Curtis CR, Erb HN, Sniffen CJ, Smith RD, Kronfeld DS. Path analysis of dry period nutrition, postpartum metabolic and reproductive disorders, and mastitis in Holstein cows. J Dairy Sci. 1985; 68:2347 -60.
- Reinhardt TA, Lippolis JD, McCluskey BJ, Goff JP, Horst RL. Prevalence of subclinical hypocalcemia in dairy herds. Vet J. 2011;188(1):122 -4.
- DeGaris PJ, Lean IJ. Milk fever in dairy cows: A review of pathophysiology and control principles. Vet J. 2008;176(1):58 -69
- Goff JP. The monitoring, prevention, and treatment of milk fever and subclinical hypocalcemia in dairy cows. Vet J. 2008;176(1):50 -7.
- Kimura K, Reinhardt TA, Goff JP. Parturition and hypocalcemia blunts calcium signals in immune cells of dairy cattle. J Dairy Sci. 2006; 89:2588 -95.
- Alonso Diez AJ. (1997). Prophylaxis de la paresia puerperal hipocalcémica bovina Med Vet 11:610-614
- Bednarek D, Kondracki M, Gzeda M. (2000). Magnesium and Calcium in prophylaxis of parturient paresis in dairy cows. Abstracts. Medycyna Weterynaryjna 56:367-371.
- Conterras P. (2002a). Hipomagnesemia: efectos y procedimientos de prevención en los rebaños. VII Congreso de la Sociedad Española de Medicina Interna Veterinaria. León: Universidad de León: 20-29.
- Conterras PA. (2002b). Paresia puerperal hipocalcémica. VII Congreso de la Sociedad Española de Medicina Interna Veterinaria. León. Universidad de León :30-34.

Intravenous administration.

WITHDRAWAL PERIOD

Zero days.

ADVERSE REACTIONS

In very rare cases, heart rhythm disturbances such as arrhythmias, tachycardia/ bradycardia and even cardiac arrest may occur during intravenous administration. Respiratory disturbances, tremors, collapse and death may also occur. If these alterations are observed, treatment should be discontinued.

SPECIAL PRECAUTIONS FOR STORAGE

This veterinary medication does not require special storage conditions.

PRESENTATIONS

100ml and 500 ml vials.

Registry no. 3948 ESP

Medication subject to veterinary prescription.

Administration under veterinary control or supervision.

