

With
carob gum



Farmalac®
Dietetic Complementary Feed



Indicated in neonatal diarrhoea

- ✓ Corrects dehydration
- ✓ Improves the viability of calves with mechanical gastrointestinal disorders

Balanced formula

- ✓ With milk and nutritional supplements
- ✓ Contains carob gum, an antidiarrheal of natural origin
- ✓ Contains extract of *Saccharomyces cerevisiae*
- ✓ High palatability

Excellent results

- ✓ Very effective in rearing replacement heifers

LABIANA
always works



Farmalac double action

Carob gum:

It acts as a mechanical protection of the intestinal villi, preventing the pathogenic bacteria from adhering to them.

Pathogenic bacteria

Carob gum

Intestinal villi

Extract of *Saccharomyces cerevisiae*:

It acts as a prebiotic, promoting the growth of the intestinal flora that competes with the pathogenic bacteria in affinity for enterocytes.

Pathogenic bacteria

Lactobacillus

Extract of *S. cerevisiae*

Intestinal villi

COMPOSITION

Dextrose, skimmed milk and whey, pectin, sodium bicarbonate, cellular wall extract of *Saccharomyces cerevisiae*, magnesium chloride, potassium chloride, sodium chloride and calcium chloride.

Additives per kg:

Preservatives:

Sodium diacetate (E 262)..... 6 g

Emulgent agents:

Carob gum (E-410)50 g

Vitamins, provitamins and chemically well-defined substances of similar effect:

Vitamin A / Retinyl palmitate (3a672b) 1,100,000 IU

Vitamin D3 (E671) 210,000 IU

Vitamin C / Ascorbic acid (3a300).....1.15 g

Vitamin B12 / Cyanocobalamin525 mg

Calcium D-Pantothenate (3a841).....113.04 mg

Vitamin B2 / Riboflavin.....104 mg

Vitamin E / All-rac- α -tocopheryl acetate (3a700)..... 101 mg

Niacinamide (3a315) 100.5 mg

Vitamin B1 / Thiamine mononitrate (3a821)60 mg

Vitamin K3 / Menadione sodium bisulfite (3a710)..... 46.3 mg

Vitamin B6 / Pyridoxine HCl (3a831)..... 37.5mg

INDICATIONS

Prevention and treatment of digestive disturbance (diarrhoea) and its recovery.

Particular nutritional purpose: stabilisation of water and electrolyte balance to support the physiological digestion.

DOSAGE AND ADMINISTRATION

Oral, in drinking water or milk.

Dissolve in drinking water or in skimmed milk at 37 °C.

Calves: 50-100 g/1.5 litres of milk or water, twice a day for 3-5 days.

Lambs and kids: 20 g/300 ml of milk or water, twice a day for 3-5 days.

Recommended length of use from 1 to 7 days.

To be used up to 16 weeks of life of the animal, at maximum.

It is recommended that advice from a veterinarian be sought before use and before extending the period of use.

PRECAUTIONS AND ADVERSE REACTIONS

Do not administer together with "antibiotics", "growth factors" and "coccidiostats".

The accidental intake of large quantities of the product in its concentrated form can cause gastric disturbances and nervous disorders. Follow your veterinary's instructions.

SPECIAL PRECAUTIONS FOR STORAGE

Store in a cool, dry place, below 25 °C, protected from light

PRESENTATIONS

100 g sachet.

1 kg bag.

αESP-08100341