

Treatment of infections in horses

 Treatment of infections of the lower respiratory tract in horses caused by aerobic Gramnegative bacteria susceptible to gentamicin

Excellent absorption and fast-acting

 Reaches maximum concentration in 30-60 minutes


The best-selling gentamicin in Spain

✓ With a good quality/price ratio





Mechanism of action of gentamicin



COMPOSITION PER ML

INDICATIONS

For the treatment of infections of the lower respiratory tract in horses caused by aerobic Gramnegative bacteria susceptible to gentamicin.

DOSAGE AND ROUTE OF ADMINISTRATION

Horses: single dose of 6600 I.U./kg l.w. once daily (equivalent to 8.25 ml/ 100 kg l.w.) for 3–5 consecutive days. Intravenous route.

WITHDRAWAL PERIOD

Horses:

Not authorized for use in horses producing meat or milk for human consumption.

CONTRAINDICATIONS

- Do not use in known cases of kidney dysfunction.
- Do not use in case of hypersensitivity to the active substance or to any excipient.
- Do not exceed the recommended dosage.
- Do not use during pregnancy.
- Do not use in dehydrated animals or with hearing alteration (loss of balance or inner ear damaged).
- Use of gentamicin on foals and neonates is not recommended.

PRESENTATIONS

100 and 250 ml vials.

Registry no. 638 ESP

Medication subject to veterinary prescription.

Administration under control or supervision of the veterinary.

