



+
MORE
Ketoprofen

15%

-
LESS
withdrawal
period

Labiprofen[®]
150 mg/ml

LABIANA
always works















Administration under control or supervision of the veterinary.

Labiprofen[®] 150 mg/ml **Injectable solution**



Benefits of a NSAID with higher concentration of Ketoprofen

KETOPROFEN		15%			VS.	10%		
DOSAGE								
		1 ml/50 kg b.w / day	1 ml/50 kg b.w / day	0,75 ml/50 kg b.w / day		1 ml/33 kg b.w / day	1 ml/33 kg b.w / day	1 ml/45 kg b.w / day
WITHDRAWAL PERIOD								
	Meat and offal	2 days	3 days	1 days		4 days	4 days	4 days
	Milk	0 hours	-	Not authorized		0 hours	-	Not authorized

COMPOSITION PER ML

Ketoprofen..... 150 mg
Benzyl alcohol10 mg

INDICATIONS

Cattle:

- Reduction of inflammation and post-partum pain, musculoskeletal disorders and lameness.
- Reduction of fever associated with bovine respiratory disease in combination with antimicrobial therapy where appropriate.
- Reduction of inflammation, fever and pain in acute clinical mastitis in combination with antimicrobial therapy where appropriate.

Pigs:

- Reduction of pyrexia in cases of respiratory disease and Postpartum Dysglactia Syndrome PDS- (Mastitis, Metritis, Agalactia syndrome) in sows, in combination with antimicrobial therapy, where appropriate.

Horses:

- Reduction of inflammation and pain associated with osteoarticular and musculoskeletal disorders (lameness, laminitis, osteoarthritis, synovitis, tendinitis, etc.).
- Reduction of postoperative pain and inflammation.
- Reduction of visceral pain associated with colic.

DOSAGE AND ROUTE OF ADMINISTRATION

- Cattle:** 1 ml of Ketoprofen per 50 kg of b.w./day for 1-3 days via IV or IM, preferably in the neck region.
- Pig:** 1 ml of Ketoprofen per 50 kg of b.w./day via IM in a single occasion. Treatment may be repeated at intervals of 24-hours during a maximum of three treatments. Each injection should be given at a different site.
- Horses:** 0.75 ml of medicine per 50 kg of b.w./day via IV for 1-5 days. In the case of colic one injection is normally sufficient.

WITHDRAWAL PERIOD

Meat:

Cattle: 2 days
Pig: 3 days
Horses: 1 day

Milk:

Cattle: Zero hours.

PRECAUTIONS AND ADVERSE REACTIONS

Do not use in animals where there is the possibility of gastro-intestinal ulceration or bleeding, in order not to aggravate their situation.

Do not use in animals suffering from cardiac, hepatic, or renal disease.

Do not use in cases of known hypersensitivity to ketoprofen or acetyl-salicylic acid or to any of the excipients.

Do not use in animals with evidence of blood dyscrasia or coagulopathy. Do not administer other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other.

SPECIAL PRECAUTIONS FOR USE

Do not exceed the recommended dose. Do not exceed the recommended treatment period.

The use of ketoprofen is not recommended in foals less than one month of age.

When administering to animals of less than 6 weeks of age, ponies or in aged animals it is necessary to adjust the dose accurately as well as to perform a close clinical follow-up.

Avoid intra-arterial injection.

Avoid use in any dehydrated, hypovolaemic or hypotensive animals as there is a potential risk of increased renal toxicity.

The use of ketoprofen is not recommended in pigs affected by PMWS (multisystemic post-weaning weight loss syndrome), in order not to aggravate their condition.

In horses, avoid extravascular administration.

USE DURING PREGNANCY, LACTATION OR LAY

Can be used in pregnant and lactating cows and lactating sows.

The use is not recommended in lactating mares.

SPECIAL PRECAUTIONS FOR STORAGE

Keep the vial in the outer carton in order to protect from light.

PRESENTATION

50, 100 and 250 ml vials

Registry no. 3966 ESP

Medication subject to veterinary prescription.

Administration under control or supervision of the veterinary.

