



A single
injection



Oxilab Ciclina Retard 200 mg/ml

Oxytetracycline in injectable solution

**The highest rated by the
veterinarians in Spain**

- ✓ Wide range of indications
- ✓ Considered in the market as
a Premium oxytetracycline

**The difference lies
in its formulation**

- ✓ Painless application
- ✓ Formulated to reduce reaction
at the point of inoculation
- ✓ Excellent tissue absorption

**Registered
in caprine livestock**




LABIANA
always works

Oxylabibiclina Retard 200 mg/ml **Injectable solution**



Mechanism of action of oxytetracycline



COMPOSITION PER ML

Oxytetracycline (dihydrate)200 mg

INDICATIONS

Treatment of de infections caused by microorganisms sensitive to oxytetracycline in bovine, ovine and caprine livestock.

Bovine:

- Respiratory infections caused by *Histophilus somni*, *Mannheimia haemolytica*, *Pasteurella multocida* and *Mycoplasma bovis*.
- Genital infections caused by *Arcanobacterium pyogenes*.
- Foot infections caused by *Dichelobacter nodosus*, *Fusobacterium necrophorum*, *Prevotella melaninogenicus*.

Ovine and caprine:

- Respiratory infections caused by *Mannheimia haemolytica*, *Pasteurella multocida*.
- Genital infections caused by *Arcanobacterium pyogenes*, *Chlamydia abortus*.
- Foot infections caused by *Dermatophilus congolensis*, *Dichelobacter nodosus*, *Fusobacterium necrophorum*, *Prevotella melaninogenicus*.

Other infections:

- Mastitis caused by *Staphylococcus aureus*, *Streptococcus agalactiae*, *Escherichia coli*.
- Enzootic abortion caused by *Chlamydia abortus*, *Chlamydia psittaci*.
- Polyarthritis caused by *Chlamydia* spp. and *Mycoplasma* spp.

DOSAGE AND ROUTE OF ADMINISTRATION

Bovine, ovine and caprine 20 mg of oxytetracycline/kg live weight in single dose (equivalent to 1 ml/10 kg l.w.) deep intramuscular route.

Intramuscular injection will be administered preferably in the muscle of the neck, except in ovine and caprine, species in which the product must be injected into the muscle of the hip zone.

Respect a maximum volume per injection point of:

Bovine: 10 ml.

Ovine and caprine: 3 ml.

Anticipate a sufficient separation between injection points when several places of administration are necessary.

WITHDRAWAL PERIOD

Bovine:

Meat: 22 days. **Milk:** 9 days (216 hours).

Ovine and caprine:

Meat: 20 days. **Milk:** Do not administer to animals in lactation whose milk is used for human consumption.

CONTRAINDICATIONS

- Do not use in case of known hypersensitivity to tetracyclines or to any excipient.
- Do not use in animals with liver or kidney disorders.
- Do not use in horses, dogs or cats.

PRECAUTIONS AND ADVERSE REACTIONS

- Local transitory reactions can appear at the point of injection (swelling, pain).
- As with the other tetracyclines, allergic and photosensitivity reactions can appear.
- In young animals, it can lead to atypical colouring of bones and teeth due to the depositing of oxytetracycline.

SPECIAL PRECAUTIONS FOR STORAGE

Shelf life: 3 years stored in its original closed bottle.

Period of validity after opening the primary container: 28 days.

PRESENTATIONS

100 and 250 ml vials.

Registry no. 2688 ESP

Medication subject to veterinary prescription.

Administration under control or supervision of the veterinary.

